

Application of the ToxTracker reporter assay in a mode of action approach for genetic toxicology assessment

Remco S. Derr¹, Peter I. Racz¹, Lisette Zeijdel¹, Harry Vrieling² and Giel Hendriks¹

Differential responses of the Bscl2

and Rtkn-GFP reporters allows

discrimination between clastogeni

ToxTracker reporter cell lines were

exposed to a selection of microtubule

disrupting agents. Induction levels of

equitoxic concentration that induced

50% cytotoxicity. The kinetics of

Bscl2-GFP and Rtkn-GFP reporter

induction was determined following

(cisplatin, etoposide, mitomycin

doxorubicin) or mitotic spindle poisons

taxol). GFP induction was determined

after 4, 8, 12, 16 and 24 h. exposure.

Exposure times that resulted in a

1.5-fold increase in GFP signal for the

Bscl2-GFP and Rtkn-GFP reporters

after exposure to various clastogenic

and aneugenic compounds were

calculated by linear regression of the

exposure time data points of the two

induction data points. The dashed lines

indicat the thresholds for classification

as clastogen or aneugen.

¹Toxys B.V., Robert Boyleweg 4, 2333 CG, Leiden, the Netherlands. g.hendriks@toxys.com. ²Department of Human Genetics, Leiden University Medical Center, Leiden, the Netherlands

Introduction

ToxTracker is a mammalian stem cell-based reporter assay that detects activation of DNA damage, oxidative stress and protein damage in a single test (Hendriks et al., Tox Sci 2016). ToxTracker can be particularly useful in an Adverse Outcome Pathway (AOP) approach for both genotoxic and non-genotoxic carcinogens.

Results

In an extensive validation study using 300 reference chemicals and >200 proprietary compounds, ToxTracker was able to discriminate between a mutagenic and classified the genotoxic compounds with a sensitivity of 94% and specificity of 95%. By assessing the differential induction of the two DNA damage reporters, ToxTracker was able to discriminate between a mutagenic and classified the genotoxic compounds with a sensitivity of 95%. By assessing the differential induction of the two DNA damage reporters, ToxTracker was able to discriminate between a mutagenic and classified the genotoxic compounds with a sensitivity of 95%. By assessing the differential induction of the two DNA damage reporters, ToxTracker was able to discriminate between a mutagenic and classified the genotoxic compounds with a sensitivity of 95%. By assessing the differential induction of the two DNA damage reporters, ToxTracker was able to discriminate between a mutagenic and classified the genotoxic compounds with a sensitivity of 95%. By assessing the differential induction of the two DNA damage reporters, ToxTracker was able to discriminate between a mutagenic and classified the genotoxic compounds with a sensitivity of 95%. By assessing the differential induction of the two DNA damage reporters, and classified the genotoxic compounds with a sensitivity of 95%. By assessing the differential induction of the two DNA damage reporters, and classified the genotoxic compounds with a sensitivity of 95%. could discriminate between a clastogenic and aneugenic mode of action by the selective induction of this reporter cell lines, ToxTracker can identify an aneugenic MOA by inhibition of cell cycle kinases.


Conclusion

§The integrative approach of the ToxTracker assay provides a unique tool for in vitro carcinogenic hazard identification of specific cellular signalling pathways upon exposure and deliver insight into the underlying mechanism of toxicity.

The ToxTracker reporter assay


- Stem cell-based reporter assay
- In vitro carcinogenicty hazard screening
- Insight into mechanisms of genotoxicity
- · Six independent GFP reporter cell lines
- High throughput detection by flow cytometry
- · Discriminate between induction of DNA damage, oxidative stress and protein damage


Biomarker gene	Mode of action	Cellular pathway
Bscl2	Mutagenic DNA lesions	ATR/Chk1 DNA damage signaling
Rtkn	DNA double-strand breaks	NF-kB signaling
Srxn1	ROS production	Nrf2 antioxidant response
Blvrb	ROS production	Nrf2-independent
Ddit3	Protein damage	Unfolded protein response
Btg2	Cytotoxicity	p53 signaling





Test name	Sensitivity (%)	Specificity (%)
Regulatory		
Bacterial reversion (Ames)	60	77
Chromosome aberrations	70	55
Mammalian mutation	81	48
Screening		
ToxTracker	94	95
Ames MPF	58	63
GreenScreen HC	87	95

Selective activation of the ToxTracker reporter cell lines. GFP reporter cells concentrations of the DNA damaging agents cisplatin and etoposide, the oxidative stress-inducing agents diethyl maleate and sodium arsenite, and the UPR-activating compounds tunicamycin and nitrophenol. GFP induction levels in intact cells were determined by flow 50 100 150 200 250 cytometry at 24 h. after initiation of the 0 0.5 1 1.5 exposure. The relative cell survival was determined by cell count after 24 h. Conc. (ug/ml) exposure using flow cytometry. Sodium arsenite 200 400 600 800 1000 Conc. (uM) Conc. (uM) Conc. (uM)









of phospho-H3 levels and polyploidy are considered hallmarks of inhibition of cell cycle kinases. The Aurora kinase inhibitors AMG900 and VX680 strongly reduced phospo-H3 levels and induced polyploidy. In contrast,

nocodazole arrested cells in mitosis and therefore increase phosho-H3 levels. The DNA damaging agents cisplatin and etoposide did not influence phosphorylation of H3 or DNA content of the cells.