




## cAMP NOMAD-FP650 CELL LINES


### -GLUCAGON-LIKE PEPTIDE 1 RECEPTOR (GLP1R)-



Red <sub>cAMP</sub>Nomad-GLP1R (HEK293 cell line)

EC<sub>50</sub> GLP-1: 2.40x10<sup>-10</sup> M

Z': 0.68+/- 0.01



**Product Name:** GLP1R <sub>cAMP</sub>Nomad cell line

**Reference:** P70503

**Recp. Official Full Name:** Glucagon-like peptide 1 receptor

**DNA Accession Number:** NM\_002062.4

**Host Cell:** HEK293

**Resistance:** G418 + Puromycin

**Quantity:** > 3 x 10<sup>6</sup> cells / vial

**Storage:** Liquid Nitrogen

### **Assay Briefly description**

Each vial of red <sub>cAMP</sub>Nomad-GLP1R contains HEK293 cells stably expressing red <sub>cAMP</sub>Nomad biosensor and Glucagon-like peptide 1 receptor (with no tag).

Innoprot's red <sub>cAMP</sub>Nomad-GLP1R cell line has been designed to assay compounds or analyze their capability to modulate Glucagon-like peptide 1 receptor. When an agonist binds to GLP1R a G protein is activated, which in turn, triggers a cellular response mediated by cAMP.


This cell line has been validated measuring cAMP increase in the cytosol analyzing red <sub>cAMP</sub>Nomad biosensor distribution within the cell. This cell line allows the image analysis of the stimuli induced by the compounds.

This highly reproducible assay has been validated using human GLP-1 as agonist in a High Content Analysis (HCA) and a High Throughput Analysis (HTA).

### **About Red <sub>cAMP</sub>Nomad Biosensor**

Red <sub>cAMP</sub>Nomad Biosensor is a fluorescent polypeptide that in the presence or absence of cAMP changes its localization within the cell.

Before cAMP production stimulation, the fluorescent biosensor is localized in the cellular membrane. An increase in this second messenger concentration leads to a change in the structural folding of red <sub>cAMP</sub>Nomad Biosensor promoting its cellular relocation in the vesicular trafficking of the cells and an increase in the fluorescence intensity.



In a cell line co-expressing red <sub>cAMP</sub>Nomad Biosensor and a GPCR of interest, the activity can be easily quantified on living cells by an image analysis platform or a fluorescence plate reader.

## cAMP Assay

cAMP<sup>Nomad</sup> HEK293 cells, stably expressing Glucagon-like peptide 1 receptor (GLP1R), were stimulated with 8 log dilution series ranging from 0 to 1  $\mu$ M of GLP-1 during 24h (n=5). % Activity was calculated relative to positive (1 $\mu$ M).

### Fluorescence intensity analysis




Fig1. Red cAMP<sup>Nomad</sup> biosensor negative control and upon 1 $\mu$ M GLP-1 stimulation.

The increase in the fluorescence was detected and analyzed using “Synergy 2” microplate reader from Biotek. The EC<sub>50</sub> for GLP-1 was  $2.4 \times 10^{-10}$  M after a treatment of 24 h with the agonist. The assay was validated with an average of  $Z' = 0.68 \pm 0.01$ .




Fig2. Concentration response curve for GLP-1 in Red cAMP<sup>Nomad</sup>-GLP1R cell line analyzed using a microplate reader.